ACCORD

Cultivation advice

Purpose: French fry / Table
Early suitable for processing

Characteristics:

Maturity	7.5
Colour of skin	LG
Color of flesh	4.5
Shape of tuber	RO/O
Depth of the eyes	7.5
Regularity shape of tuber	8
Size of tubers	8
Grading	8
Dry matter percentage	21
Suitability for cooking	7BC/C
Dormancy	5

Comments:
9 = positive .... 1 = negative
Yellowness of flesh: 8 = yellow . . . . 4 = white
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
Shape of tuber: R = Round O = Oval L = Long
Cooking type: A = Firm . . . . D = Very mealy

Planting distance:

Size	Rowspace 75 cm	Plants/hectare (x 1000)
28/35	25	53,3
35/50	34	39,2
35/55	37	36,0
50/55	38	35,1
50/60	40	33,3

Fertilisation (kg/ha):

depending on the soil reserve



				Total
Nitrogen:	(N)	230		230
Phosphate:	(P_2O_5)	110		110
Potassium:	(K_2O)	230	100	330

Storage:



7 °C
4 months

Start late blight control in time
Wait for a set skin before harvest
Very sensitive for greening
Don't de-sprout the seeds before planting

Resistance:

Leaf roll	7
A-virus	8.5
X-virus	9.5
Y-virus	8
Yntn-virus	8
Foliage blight	3
Tuber blight	7
Wart disease	wv1
Potato cyst nematode	Ro1,2,3,4,5
Common scab	6
Spraing	6
Bruising	5.5
Secondary growth	7

Comments:
R = resistant .... 1 = very susceptible
Ro(x) = Globodera Rostochiensis pathotype (x)
Pa(x) = Globodera Pallida pathotype (x)
(wv)O1 = (little susceptible) wart disease fysio 1

C. Meijer B.V. will not be responsible for possible negative effects originating from this advice.