

CUPIDO

Cultivation advice



**Purpose: Tablestock
Large tubers and early**

Characteristics:

Maturity	8.0
Colour of skin	LG
Color of flesh	6.5
Shape of tuber	O
Depth of the eyes	9
Regularity shape of tuber	8
Size of tubers	7.5
Grading	8
Dry matter percentage	20
Suitability for cooking	7AB
Dormancy	8


Comments:
9 = positive 1 = negative
Yellowness of flesh: 8 = yellow 4 = white
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
Shape of tuber: R = Round O = Oval L = Long
Cooking type: A = Firm D = Very mealy

Planting distance:


Size	Rowspace 75 cm	Plants/hectare (x 1000)
28/35	18	74,1
35/45	25	53,3
35/50	27	49,4
35/55	28	47,6
45/50	31	43,0
50/55	33	40,4

Fertilisation (kg/ha):

depending on the soil reserve



Storage:



4 °C
7 months

Resistance:

Leaf roll	6
A-virus	9
X-virus	9.5
Y-virus	9.5
Yntn-virus	R
Foliage blight	2.5
Tuber blight	6
Wart disease	O1
Potato cyst nematode	Ro1
Common scab	5
Spraing	8
Bruising	6.5
Secondary growth	7

Comments:
R = resistant 1 = very susceptible
Ro(x) = Globodera Rostochiensis pathotype (x)
Pa(x) = Globodera Pallida pathotype (x)
(ww)O1 = (little susceptible) wart disease fysio 1

Do not plant early in cold soil
Avoid too strong germination
Wait for a set skin before harvest
After harvest, dry the potatoes by ventilation