FAMBO

Cultivation advice

Purpose: French fry / Table
Good processing quality

Characteristics:

Maturity	7.5
Colour of skin	DG
Color of flesh	6
Shape of tuber	O
Depth of the eyes	7
Regularity shape of tuber	8
Size of tubers	8
Grading	7
Dry matter percentage	20.5
Suitability for cooking	6BC
Dormancy	6.5

Comments:
9 = positive .... 1 = negative
Yellowness of flesh: 8 = yellow . . . . 4 = white
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
Shape of tuber: R = Round O = Oval L = Long
Cooking type: A = Firm . . . . D = Very mealy

Planting distance:

Size	Rowspace 75 cm	Plants/hectare (x 1000)
28/35	21	63,5
35/50	32	41,7
35/55	34	39,2
35/45	30	44,4
45/50	35	38,1

Fertilisation (kg/ha):

depending on the soil reserve



				Total
Nitrogen:	(N)	200	50	250
Phosphate:	(P_2O_5)	110		110
Potassium:	(K_2O)	230		230

Storage:



7 °C
4 months

Resistance:

Leaf roll	5.5
A-virus	R
X-virus	6
Y-virus	9
Yntn-virus	5
Foliage blight	3.5
Tuber blight	8
Wart disease	O1
Potato cyst nematode	-
Common scab	7
Spraing	7
Bruising	7
Secondary growth	7

Comments:
R = resistant .... 1 = very susceptible
Ro(x) = Globodera Rostochiensis pathotype (x)
Pa(x) = Globodera Pallida pathotype (x)
(wv)O1 = (little susceptible) wart disease fysio 1

Do not use Sencor
Pre-sprout the seeds well
Avoid Rhizoctonia
Use a widely shaped ridge

C. Meijer B.V. will not be responsible for possible negative effects originating from this advice.