

# LADY AMARILLA

## Cultivation advice



Purpose: French fries

### Characteristics:

Maturity	7.5
Colour of skin	DG
Color of flesh	7.5
Shape of tuber	LO/O
Depth of the eyes	8
Regularity shape of tuber	7.5
Size of tubers	7.5
Grading	7.5
Dry matter percentage	22.5
Suitability for cooking	7BC
Dormancy	8


Comments:  
 9 = positive .... 1 = negative  
 Yellowness of flesh: 8 = yellow . . . . . 4 = white  
 Colour of skin: L=light, D=dark, W=white, G=yellow, R=red  
 Shape of tuber: R = Round O = Oval L = Long  
 Cooking type: A = Firm . . . . . D = Very mealy

### Planting distance:

Size	Rowspace 75 cm	Plants/hectare ( x 1000)
28/35	31	43,0
35/50	40	33,3
35/55	42	31,7


### Fertilisation (kg/ha):

depending on the soil reserve



To prevent mineral deficiencies, apply these elements in time ( B, Mg, Mn, S )

### Storage:



5 °C  
9 months

### Resistance:

Leaf roll	7
A-virus	8
X-virus	8
Y-virus	6
Yntn-virus	7
Foliage blight	4.5
Wart disease	r1
Potato cyst nematode	-
Common scab	5.5
Spraing	8
Bruising	9
Secondary growth	8

Comments:  
 R = resistant .... 1 = very susceptible  
 Ro(x) = Globodera Rostochiensis pathotype (x)  
 Pa(x) = Globodera Pallida pathotype (x)  
 (w) r1 = (little susceptible) resistant wart disease fysio 1

Sensitive to Alternaria, start early in the season to prevent this.  
 Secure water availability at light soils