LADY BLANCA

Cultivation advice

Purpose: French fries
Early suitable for processing

Characteristics:

Maturity	7.8
Colour of skin	LR
Color of flesh	4.5
Shape of tuber	L
Depth of the eyes	8
Regularity shape of tuber	7
Size of tubers	7.5
Grading	7
Dry matter percentage	20.5
Suitability for cooking	7AB
Dormancy	6.5

Comments:
9 = positive .... 1 = negative
Yellowness of flesh: 8 = yellow . . . . 4 = white
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
Shape of tuber: R = Round O = Oval L = Long
Cooking type: A = Firm . . . . D = Very mealy

Planting distance:

Size	Rowspace 75 cm	Plants/hectare (x 1000)
28/35	30	44,4
35/45	34	39,2
35/50	36	37,0
45/50	40	33,3
50/55	42	31,7

Fertilisation (kg/ha):

depending on the soil reserve



				Total
Nitrogen:	(N)	200	50	250
Phosphate:	(P_2O_5)	110		110
Potassium:	(K_2O)	230		230

Storage:



6 °C
8 months

Resistance:

Leaf roll	7
A-virus	8
X-virus	8
Y-virus	7
Yntn-virus	9
Foliage blight	3.5
Tuber blight	7
Wart disease	O1
Potato cyst nematode	-
Common scab	5
Spraing	7
Bruising	8
Secondary growth	7.5

Comments:
R = resistant .... 1 = very susceptible
Ro(x) = Globodera Rostochiensis pathotype (x)
Pa(x) = Globodera Pallida pathotype (x)
(wv)O1 = (little susceptible) wart disease fysio 1

Be carefull by using Sencor, only little practical experience with Sencor.
To avoid green, make big ridges.
Light soils are less suitable to cultivate this variety.

C. Meijer B.V. will not be responsible for possible negative effects originating from this advice.