LADY FELICIA

Cultivation advice

Purpose: Tablestock

Characteristics:

Maturity	7.5
Colour of skin	G
Color of flesh	7.5
Shape of tuber	O
Depth of the eyes	7.5
Regularity shape of tuber	9
Size of tubers	7
Grading	9
Dry matter percentage	19.5
Suitability for cooking	8B
Dormancy	8

Comments:
9 = positive .... 1 = negative
Yellowness of flesh: 8 = yellow . . . . 4 = white
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
Shape of tuber: R = Round O = Oval L = Long
Cooking type: A = Firm . . . . D = Very mealy

Planting distance:

Size	Rowspace 75 cm	Plants/hectare (x 1000)
28/35	19	70,2
35/45	26	51,3
35/50	27	49,4
45/50	31	43,0
50/55	33	40,4
50/60	35	38,1

Fertilisation (kg/ha):

depending on the soil reserve



				Total
Nitrogen:	(N)	200	50	250
Phosphate:	(P_2O_5)	110		110
Potassium:	(K_2O)	230		230

Storage:



4 °C
7 months

Resistance:

Leaf roll	5
A-virus	9
X-virus	7
Y-virus	9
Yntn-virus	R
Foliage blight	2.5
Tuber blight	8
Wart disease	-
Potato cyst nematode	Ro1,4
Common scab	7.5
Spraing	7.5
Bruising	7
Secondary growth	8

Comments:
R = resistant .... 1 = very susceptible
Ro(x) = Globodera Rostochiensis pathotype (x)
Pa(x) = Globodera Pallida pathotype (x)
(wv)O1 = (little susceptible) wart disease fysio 1

Do not use Sencor
Wait for eyes waking up before planting
Avoid too strong germination
Start late blight control in time
Wait for a set skin before harvest
After harvest, dry the potatoes by ventilation

C. Meijer B.V. will not be responsible for possible negative effects originating from this advice.