LADY OLYMPIA

Cultivation advice

Purpose: French fry / Table
For long storage

Characteristics:

Maturity	6.5
Colour of skin	G
Color of flesh	7
Shape of tuber	LO
Depth of the eyes	7
Regularity shape of tuber	7
Size of tubers	8
Grading	7.5
Dry matter percentage	22.5
Suitability for cooking	7C
Dormancy	6.5

Comments:
9 = positive .... 1 = negative
Yellowness of flesh: 8 = yellow . . . . 4 = white
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
Shape of tuber: R = Round O = Oval L = Long
Cooking type: A = Firm . . . . D = Very mealy

Planting distance:

Size	Rowspace 75 cm	Plants/hectare (x 1000)
28/35	31	43,0
35/45	35	38,1
35/50	40	33,3
45/50	42	31,7
50/55	44	30,3

Fertilisation (kg/ha):

depending on the soil reserve



				Total
Nitrogen:	(N)	225	50	275
Phosphate:	(P_2O_5)	110		110
Potassium:	(K_2O)	230		230

Storage:



6 °C
8 months

Sensitive to the full dose of Sencor pre-emergence.
Wait for eyes waking up before planting
Start late blight control in time
Wait for a set skin before harvest
After harvest, dry the potatoes by ventilation

Resistance:

Leaf roll	7
A-virus	9
X-virus	9
Y-virus	8.5
Yntn-virus	7
Foliage blight	3
Tuber blight	5
Wart disease	O1
Potato cyst nematode	-
Common scab	6.5
Spraing	7
Bruising	7.5
Secondary growth	6

Comments:
R = resistant .... 1 = very susceptible
Ro(x) = Globodera Rostochiensis pathotype (x)
Pa(x) = Globodera Pallida pathotype (x)
(wv)O1 = (little susceptible) wart disease fysio 1

C. Meijer B.V. will not be responsible for possible negative effects originating from this advice.