

# MELODY



## Cultivation advice

**Purpose: Tablestock  
For long storage**

### Characteristics:

Maturity	6.0
Colour of skin	LG
Color of flesh	7
Shape of tuber	O
Depth of the eyes	8
Regularity shape of tuber	8
Size of tubers	7.5
Grading	8
Dry matter percentage	20.5
Suitability for cooking	8B/BC
Dormancy	8


Comments:  
9 = positive .... 1 = negative  
Yellowness of flesh: 8 = yellow . . . . 4 = white  
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red  
Shape of tuber: R = Round O = Oval L = Long  
Cooking type: A = Firm . . . . D = Very mealy

### Planting distance:


Size	Rowspace 75 cm	Plants/hectare ( x 1000)
28/35	20	66,7
35/45	26	51,3
35/50	28	47,6
35/55	30	44,4
45/50	31	43,0
50/55	33	40,4

### Fertilisation (kg/ha):

depending on the soil reserve



### Storage:



5 °C  
7 months

### Resistance:

Leaf roll	8
A-virus	8
X-virus	8.5
Y-virus	7.5
Yntn-virus	R
Foliage blight	5.5
Tuber blight	7.5
Wart disease	O1
Potato cyst nematode	Ro1
Common scab	7.5
Spraing	7
Bruising	7.5
Secondary growth	7.5

Comments:  
R = resistant .... 1 = very susceptible  
Ro(x) = Globodera Rostochiensis pathotype (x)  
Pa(x) = Globodera Pallida pathotype (x)  
(wv)O1 = (little susceptible) wart disease fysio 1

Wait for eyes waking up before planting  
After harvest, dry the potatoes by ventilation  
Avoid silver scurf  
No sprouting inhibitor during handling  
Sensitive to the full dose of Sencor pre-emergence.