MURATO

Cultivation advice

Purpose: Redskinned table potato
Export

Characteristics:

Maturity	6
Colour of skin	R
Color of flesh	6
Shape of tuber	LO
Depth of the eyes	8
Regularity shape of tuber	7
Size of tubers	7
Grading	7
Dry matter percentage	20.7
Suitability for cooking	6B
Dormancy	5

Comments:
9 = positive .... 1 = negative
Yellowness of flesh: 8 = yellow . . . . 4 = white
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
Shape of tuber: R = Round O = Oval L = Long
Cooking type: A = Firm . . . . D = Very mealy

Planting distance:

Size	Rowspace 75 cm	Plants/hectare (x 1000)
35/55	30	44,4

Fertilisation (kg/ha):

depending on the soil reserve



				Total
Nitrogen:	(N)	225	50	275
Phosphate:	(P_2O_5)	110		110
Potassium:	(K_2O)	230	100	330

Storage:



6 °C
4 months

Resistance:

Leaf roll	6
A-virus	
X-virus	
Y-virus	7
Yntn-virus	
Foliage blight	4
Tuber blight	7
Wart disease	-
Potato cyst nematode	Ro1,4
Common scab	6
Spraing	8
Bruising	5
Secondary growth	7

Comments:
R = resistant .... 1 = very susceptible
Ro(x) = Globodera Rostochiensis pathotype (x)
Pa(x) = Globodera Pallida pathotype (x)
(wv)O1 = (little susceptible) wart disease fysio 1

Do not plant early in cold soil
Avoid black spot
Do not use Sencor

C. Meijer B.V. will not be responsible for possible negative effects originating from this advice.