MUSICA

Cultivation advice

Purpose: Tablestock

Characteristics:

Maturity	7
Colour of skin	G
Color of flesh	8
Shape of tuber	L/LO
Depth of the eyes	7.5
Regularity shape of tuber	7
Size of tubers	7.5
Grading	8
Dry matter percentage	19
Suitability for cooking	8A
Dormancy	4.5

Comments:
9 = positive .... 1 = negative
Yellowness of flesh: 8 = yellow . . . . 4 = white
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
Shape of tuber: R = Round O = Oval L = Long
Cooking type: A = Firm . . . . D = Very mealy

Planting distance:

Size	Rowspace 75 cm	Plants/hectare (x 1000)
Plantdistance for cultivation size 40/70 mm		
28/35	20	66,7
35/50	28	47,6
35/55	30	44,4
50/55	33	40,4

Fertilisation (kg/ha):

depending on the soil reserve



				Total
Nitrogen:	(N)	200	50	250
Phosphate:	(P_2O_5)	110		110
Potassium:	(K_2O)	230		230

To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, S)

Storage:



4 °C
6 months

Do not pre-sprout
Heavy clay soils increase % misshapen tubers
Best skin finish before the crop gets mature naturally

Resistance:

Leaf roll	7
A-virus	7
X-virus	8
Y-virus	8
Yntn-virus	R
Foliage blight	4
Wart disease	r1
Potato cyst nematode	Ro1,2,3,Pa2
Common scab	6.5
Spraing	6.5
Bruising	8
Secondary growth	6.5

Comments:
R = resistant .... 1 = very susceptible
Ro(x) = Globodera Rostochiensis pathotype (x)
Pa(x) = Globodera Pallida pathotype (x)
(w) r1 = (little susceptible) resistant wart disease fysio 1

C. Meijer B.V. will not be responsible for possible negative effects originating from this advice.