

ORCHESTRA



Cultivation advice

Purpose: Tablestock

Characteristics:

Maturity	7.5
Colour of skin	LG
Color of flesh	6.5
Shape of tuber	RO/O
Depth of the eyes	7.5
Regularity shape of tuber	8.5
Size of tubers	8
Grading	8
Dry matter percentage	17.5
Suitability for cooking	7.5A
Dormancy	5.5


Comments:
 9 = positive 1 = negative
 Yellowness of flesh: 8 = yellow 4 = white
 Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
 Shape of tuber: R = Round O = Oval L = Long
 Cooking type: A = Firm D = Very mealy

Planting distance:


Size	Rowspace 75 cm	Plants/hectare (x 1000)
28/35	24	55,6
35/45	30	44,4
35/50	32	41,7
35/55	34	39,2
45/50	35	38,1
50/55	37	36,0

Fertilisation (kg/ha):

depending on the soil reserve



Storage:



5 °C
6 months

Resistance:

Leaf roll	9
A-virus	8
X-virus	8
Y-virus	9
Yntn-virus	R
Foliage blight	3.5
Tuber blight	
Wart disease	wvO1
Potato cyst nematode	Ro1
Common scab	7
Spraing	8
Bruising	7
Secondary growth	8

Comments:
 R = resistant 1 = very susceptible
 Ro(x) = Globodera Rostochiensis pathotype (x)
 Pa(x) = Globodera Pallida pathotype (x)
 (wv)O1 = (little susceptible) wart disease fysio 1

Do not use Sencor. Very sensitive to metribuzin.
 No sprouting inhibitor during handling