

TURBO

Cultivation advice



Purpose: Tablestock

Characteristics:

Maturity	7.0
Colour of skin	LG
Color of flesh	6
Shape of tuber	LO
Depth of the eyes	7
Regularity shape of tuber	7
Size of tubers	7.5
Grading	6.5
Dry matter percentage	21
Suitability for cooking	7BC
Dormancy	6


Comments:
 9 = positive 1 = negative
 Yellowness of flesh: 8 = yellow 4 = white
 Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
 Shape of tuber: R = Round O = Oval L = Long
 Cooking type: A = Firm D = Very mealy

Planting distance:


Size	Rowspace 75 cm	Plants/hectare (x 1000)
28/35	19	70,2
35/45	27	49,4
35/50	29	46,0
35/55	30	44,4
45/50	31	43,0
50/55	34	39,2

Fertilisation (kg/ha):

depending on the soil reserve



Storage:



5 °C
7 months

Resistance:

Leaf roll	4
A-virus	7.5
X-virus	6
Y-virus	6.5
Yntn-virus	8
Foliage blight	4.5
Tuber blight	8
Wart disease	O1
Potato cyst nematode	Ro1,2,3
Common scab	7
Spraing	5.5
Bruising	8
Secondary growth	5

Comments:
 R = resistant 1 = very susceptible
 Ro(x) = Globodera Rostochiensis pathotype (x)
 Pa(x) = Globodera Pallida pathotype (x)
 (ww)O1 = (little susceptible) wart disease fysio 1

Pre-sprout the seeds well
 Avoid Rhizoctonia
 Do not plant early in cold soil
 Sensitive to the full dose of Sencor pre-emergence.